arXiv:1601.06964v2 [physics.atom-ph] 7 Feb 2016

Subcomponents $\psi_{4,1}^{(2d)}$ and $\psi_{3,0}^{(2c)}$ of the angular Fock coefficients: Explicit analytic representation

Evgeny Z. Liverts and Nir Barnea

Racah Institute of Physics, The Hebrew University, Jerusalem 91904, Israel

Abstract

The Fock expansion [1] describes the S-state wave function of the two-electron atomic system in the vicinity of the triple coalescence point. The present work constitutes the additional appendix to our paper [2] devoted to refinement and further development of calculation of the angular Fock coefficients of the Fock expansion. We derive the explicit analytic expressions for the complicated subcomponents $\psi_{4,1}^{(2d)}$ and $\psi_{3,0}^{(2c)}$ of the angular Fock coefficients. An unusual method of using the Wolfram *Mathematica* was applied.

PACS numbers: 31.15.-p, 31.15.A-, 31.15.xj, 03.65.Ge

I. INTRODUCTION

In our paper [2] the various components of the angular Fock coefficients (AFC) were derived in the form of the one-dimensional series with fast convergence. In particular, the solution of the individual Fock recurrence relation (see Eq.(C1))

$$\left(\Lambda^2 - 32\right)\psi_{4,1}^{(2d)} = h_{4,1}^{(2d)} \tag{1}$$

was represented by the subcomponent

$$\psi_{4,1}^{(2d)}(\alpha,\theta) = \sum_{l=0}^{\infty} P_l(\cos\theta)(\sin\alpha)^l \mathrm{t}_l(\alpha), \tag{2}$$

where $P_l(z)$ is the Legendre polynomial. Treating $r_1,\ r_2$ and r_{12} as the interparticle coordinates the hyperspherical angles α and θ are defined as

$$\alpha = 2\arctan\left(r_2/r_1\right), \quad \theta = \arccos\left[(r_1^2 + r_2^2 - r_{12}^2)/(2r_1r_2)\right]. \tag{3}$$

The hyperspherical angular momentum operator reads (see Eq.(5) [2])

$$\Lambda^2 = -\frac{4}{\sin^2\alpha}\left(\frac{\partial}{\partial\alpha}\sin^2\alpha\frac{\partial}{\partial\alpha} + \frac{1}{\sin\theta}\frac{\partial}{\partial\theta}\sin\theta\frac{\partial}{\partial\theta}\right), \tag{4}$$

and the right-hand side (rhs) of Eq.(1) is (see, Eqs.(C2), (C3) [2])

$$h_{4,1}^{(2d)}(\alpha,\theta) = \sum_{l=0}^{\infty} P_l(\cos\theta)(\sin\alpha)^l \mathrm{h}_l(\alpha), \tag{5}$$

where

$$\begin{aligned}\mathrm{h}_l(\alpha) = {} & \frac{\pi-2}{3\pi(2l-1)2^l}\left[\sin\left(\frac{\alpha}{2}\right) + \cos\left(\frac{\alpha}{2}\right)\right]\times \\ & \left[\frac{5}{(2l-3)\sin\alpha}F_{l,3}(\alpha) - \left(1 - \frac{2}{\sin\alpha}\right)F_{l,1}(\alpha) - (2l-1)F_{l,-1}(\alpha)\right].\end{aligned} \tag{6}$$

Definition

$$F_{l,\nu}(\alpha) = {}_2F_1\left(\frac{l}{2} - \frac{\nu}{4}, \frac{l}{2} - \frac{\nu}{4} + \frac{1}{2}; l + \frac{3}{2}; \sin^2\alpha\right) \tag{7}$$

was used for the Gauss hypergeometric function in the rhs of Eq.(7).

It was shown in [2] that functions $\tau_l(\rho) \equiv t_l(\alpha)$ with $\rho = \tan(\alpha/2)$ represent the physical solution of equation

$$\left(1+\rho^2\right)^2\tau_l''(\rho) + 2\rho^{-1}\left[1 + \rho^2 + l(1-\rho^4)\right]\tau_l'(\rho) + 4(2-l)(l+4)\tau_l(\rho) = -h_l(\rho), \tag{8}$$

were (see Eq.(C6) [2])

$$h_l(\rho) \equiv \mathrm{h}_l(\alpha) = -\frac{(\pi-2)(\rho+1)\left(\rho^2+1\right)^{l-1}}{3\pi(2l-1)(2l+3)2^{l+1}} \times$$
$$\left[\frac{15-4l(l+1)(4l+11)}{(2l-3)(2l+5)\rho} + 4l(2l+3) + 2\rho + 4(l+1)(2l-1)\rho^2 + \frac{(2l-1)(4l+5)\rho^3}{2l+5}\right]. \quad (9)$$

The physical solutions of Eq.(8) for $l = 0, 1, 2$ were presented in [2] by Eqs.(94)-(96), respectively. Putting $A_1 = 0$ in Eq.(C17) [2], one obtains the physical solution of Eq.(8) for $l > 2$ in the form

$$\tau_l(\rho) = \tau_l^{(p)}(\rho) + A_2 v_{4l}(\rho), \qquad (l \geq 3) \quad (10)$$

where the particular solution $\tau_l^{(p)}$ of the differential equation (8) can be obtained by Eq.(45) [2] with $k = 4$. This gives

$$\tau_l^{(p)}(\rho) = \frac{1}{2l+1}\left[u_{4l}(\rho)\mathcal{V}_l(\rho) - v_{4l}(\rho)\mathcal{U}_l(\rho)\right], \quad (11)$$

where

$$\mathcal{V}_l(\rho) = \int_0^\rho v_{4l}(t)h_l(t)t^{2l+2}(1+t^2)^{-2l-3}dt, \quad (12)$$

$$\mathcal{U}_l(\rho) = \int_1^\rho u_{4l}(t)h_l(t)t^{2l+2}(1+t^2)^{-2l-3}dt. \quad (13)$$

The solutions of the homogeneous equation associated with Eq.(8) is defined by Eq.(46)[2] for $k = 4$, which yields

$$u_{4l}(\rho) = \rho^{-2l-1}(\rho^2+1)^{l+4}\ {}_2F_1\left(\frac{7}{2}, 3-l; \frac{1}{2}-l; -\rho^2\right), \quad (14a)$$

$$v_{4l}(\rho) = (\rho^2+1)^{l+4}\ {}_2F_1\left(\frac{7}{2}, 4+l; l+\frac{3}{2}; -\rho^2\right). \quad (14b)$$

II. DERIVATION OF THE EXPLICIT REPRESENTATION FOR THE PARTICULAR SOLUTION $\tau_l^{(p)}$

To solve the problem mentioned in the title of this Section, it would be useful, first of all, to express the solutions $u_{4l}(\rho)$ and $v_{4l}(\rho)$ of the homogeneous equation through the elementary functions. It is seen from Eq.(14a) that for $l \geq 3$, we can write down

$$u_{4l}(\rho) = \frac{8(l-3)!(1+\rho^2)^{l+4}}{15\sqrt{\pi}\Gamma(l+1/2)\rho^{2l+1}} \sum_{m=0}^{l-3} \frac{(-1)^m\Gamma(m+7/2)\Gamma(l-m+1/2)}{m!(l-m-3)!}\rho^{2m}. \quad (15)$$

The well-known formula (7.3.1.140)[3] was applied.

Solution of the problem for $v_{4l}(\rho)$, defined by Eq.(14b), is more complicated. Making use of the relation (7.3.1.9) [3] for $m = l - 3$ (alternatively, one can apply [4]), and subsequent application of the transformation (7.3.1.3) [4] (alternatively, one can apply [5]) yields

$$ {}_2F_1\left(\frac{7}{2}, l+4; l+\frac{3}{2}; -\rho^2\right) = \frac{16\Gamma(l+3/2)}{105\sqrt{\pi}\rho^{2(l-3)}(1+\rho^2)^7} \sum_{p=0}^{l-3} \frac{(-1)^p}{p!(l-p-3)!} \times {}_2F_1\left(1, l-p+4; \frac{9}{2}; \frac{\rho^2}{1+\rho^2}\right). \quad (16) $$

Next step is application of the relation (7.3.1.132) [3] or the Wolfram Function Site [6] to the Gauss Hypergeometric Functions in the rhs of Eq.(16). This gives

$$ {}_2F_1\left(1, l-p+4; \frac{9}{2}; \frac{\rho^2}{1+\rho^2}\right) = \frac{7\Gamma(l-p+1/2)}{2(l-p+3)!}\left\{\frac{15(1+\rho^2)^{l-p+4}}{8\rho^8} \times \left[\frac{2\rho}{\sqrt{\pi}}\arctan(\rho) - \sum_{m=1}^{3}\frac{(m-1)!}{\Gamma(m+1/2)}\left(\frac{\rho^2}{1+\rho^2}\right)^m\right] + \sum_{m=0}^{l-p-1}\frac{(l-p-m+2)!}{\Gamma(l-p-m+1/2)}(1+\rho^2)^{m+1}\right\}. \quad (17) $$

Inserting Eqs.(16)-(17) into Eq.(14b), one obtains the following representation for the second solution of the homogeneous equation:

$$ v_{4l}(\rho) = \frac{\Gamma(l+3/2)}{\sqrt{\pi}}\left(\frac{1+\rho^2}{\rho^2}\right)^{l-3}\sum_{p=0}^{l-3}\frac{(-1)^p\Gamma(l-p+1/2)}{p!(l-p-3)!(l-p+3)!}\left\{\frac{(1+\rho^2)^{l-p+4}}{\rho^8} \times \left[\frac{2\rho}{\sqrt{\pi}}\arctan(\rho) - \sum_{m=1}^{3}\frac{(m-1)!}{\Gamma(m+1/2)}\left(\frac{\rho^2}{1+\rho^2}\right)^m\right] + \frac{8}{15}\sum_{m=0}^{l-p-1}\frac{(l-p-m+2)!}{\Gamma(l-p-m+1/2)}(1+\rho^2)^{m+1}\right\}. \quad (18) $$

To derive the analytic representation for the particular solution $\tau_l^{(p)}$, one needs to find analytic representations for the integrals (12) and (13) included into the rhs of Eq.(11). To this end, first of all, let us present the rhs of Eq.(8) (defined by Eq.(9)) in the compact form

$$ h_l(t) = a_{0l}(t+1)(t^2+1)^{l-1}\sum_{n=1}^{5} a_{nl}t^{n-2}, \quad (19) $$

where

$$ a_{0l} = -\frac{(\pi-2)2^{-l-1}}{3\pi(2l-1)(2l+3)}, \quad a_{1l} = \frac{15-4l(l+1)(4l+11)}{(2l-3)(2l+5)}, \quad a_{2l} = 4l(2l+3), $$
$$ a_{3l} = 2, \quad a_{4l} = 4(l+1)(2l-1), \quad a_{5l} = \frac{(2l-1)(4l+5)}{2l+5}. \quad (20) $$

Inserting representations (15) and (19) into the rhs of Eq.(13), and performing the trivial

integration, one obtains

$$\mathcal{U}_l(\rho) = a_{0l}\frac{8(l-3)!}{15\sqrt{\pi}\Gamma(l+1/2)}\sum_{m=0}^{l-3}\frac{\Gamma(m+7/2)\Gamma(l-m+1/2)(-1)^m}{m!(l-m-3)!}\times \\ \sum_{n=1}^{5}a_{nl}\left(\frac{\rho^{2m+n}-1}{2m+n}+\frac{\rho^{2m+n+1}-1}{2m+n+1}\right). \tag{21}$$

Using Eq.(14b) and Eqs.(19)-(20) it is convenient to present Eq.(12) in the form

$$\mathcal{V}_l(\rho) = a_{0l}\sum_{n=0}^{5}b_{n,l}\int_0^\rho t^{2l+n+1}\ {}_2F_1\left(\frac{7}{2},l+4;l+\frac{3}{2};-t^2\right)dt, \tag{22}$$

where

$$b_{0,l}=a_{1l},\quad b_{5,l}=a_{5l},\quad b_{j,l}=a_{jl}+a_{j+1l}\quad (j=1,2,3,4). \tag{23}$$

Making use of the relation (1.16.1) [3] or the the Wolfram Functions Site [7] with $a=7/2, b=l+4, c=l+3/2, \alpha=l+(p+1)/2, z=t^2$ yields

$$\mathcal{V}_l(\rho) = a_{0l}\sum_{n=0}^{5}\left(\frac{b_{n,l}}{2l+n+2}\right)\rho^{2l+n+2}\ {}_3F_2\left(\frac{7}{2},l+4,l+1+\frac{n}{2};l+\frac{3}{2},l+2+\frac{n}{2};-\rho^2\right). \tag{24}$$

The latter relation gives the representation of the integral (12) through the generalized hypergeometric functions.

It can be useful to obtain the representation of the integral (12) through the Gauss hypergeometric functions. To this end, one can apply the relation (7.4.1.2) [3] or the Wolfram Functions Site [8]. A simple subsequent reorganization of summation along with application of the linear transformation (7.3.1.4) [3] gives finally

$$\mathcal{V}_l(\rho) = \frac{4a_{0l}\Gamma(l+3/2)}{15\sqrt{\pi}}\rho^{2l+3}\sum_{m=0}^{2}\frac{m!}{(1+\rho^2)^{5-m}}\sum_{k=0}^{m}\frac{\Gamma(k+7/2)}{k!(m-k)!\Gamma(k+l+3/2)}\left(-\frac{\rho^2}{1+\rho^2}\right)^k\times \\ \left[\frac{(k+l+3)!\Gamma(l+m+3/2)}{(l+3)!\Gamma(k+l+m+5/2)}b_{2m+1,l}\rho^{2m}\ {}_2F_1\left(l+m-1,m-\frac{3}{2};k+l+m+\frac{5}{2};-\rho^2\right)\right. \\ \left.+\frac{b_{2(2-m),l}}{k+l-m+3}\rho^{3-2m}\ {}_2F_1\left(l-2,m-\frac{3}{2};k+l+\frac{3}{2};-\rho^2\right)\right]. \tag{25}$$

Now we shall show that the Gauss hypergeometric functions included into Eq.(25), in its turn, can be expressed through the elementary functions. Use of the relation (7.3.1.10) [3] or the Wolfram Functions Site [9] with $n=l-3$, and subsequent application of the linear

transformation (7.3.1.3) [3] gives

$$ {}_2F_1\left(l-2, m-\frac{3}{2}; k+l+\frac{3}{2}; -\rho^2\right) = \frac{(-\rho^2)^{3-l}\left(k+\frac{9}{2}\right)_{l-3}}{1+\rho^2} \times \sum_{p=0}^{l-3} \frac{(-1)^p}{p!(l-p-3)!}\, {}_2F_1\left(1, k+p+6-m; k+\frac{9}{2}; \frac{\rho^2}{1+\rho^2}\right). \quad (26) $$

In its turn, using (7.3.1.10) [3] or the Wolfram Functions Site [9] with $n = l+m-2$, and then applying (7.3.1.3) [3], one obtains

$$ {}_2F_1\left(l+m-1, m-\frac{3}{2}; k+l+m+\frac{5}{2}; -\rho^2\right) = \frac{\left(k+\frac{9}{2}\right)_{l+m-2}}{(1+\rho^2)\,(-\rho^2)^{l+m-2}} \times \sum_{p=0}^{l+m-2} \frac{(-1)^p}{p!(l+m-p-2)!}\, {}_2F_1\left(1, k+p+6-m; k+\frac{9}{2}; \frac{\rho^2}{1+\rho^2}\right). \quad (27) $$

Finally, application of the relation (7.3.1.132) [3] or the Wolfram Functions Site [6] yields:

$$ {}_2F_1\left(1, k+p+6-m; k+\frac{9}{2}; \frac{\rho^2}{1+\rho^2}\right) = \frac{\Gamma(k+9/2)\Gamma(p-m+5/2)(1+\rho^2)^{k-m+p+6}}{\pi(k-m+p+5)!\rho^{2(k+4)}} \times \left[2\rho\arctan(\rho) - \sqrt{\pi}\sum_{s=1}^{k+3}\frac{(s-1)!}{\Gamma(s+1/2)}\left(\frac{\rho^2}{1+\rho^2}\right)^s\right] + \frac{2k+7}{2(k-m+p+5)}\sum_{s=0}^{p-m+1}\frac{(-1)^s\left(m-p-\frac{3}{2}\right)_s(1+\rho^2)^{s+1}}{(k-m+p-s+5)_s}, \quad (28) $$

where $(a)_n$ is the Pochhammer symbol. Thus, Eqs.(25)-(28) give the representation of the integral (12) through the rational functions and the arctangent of ρ. Finally, Eqs.(25)-(28) together with Eqs.(15), (18) and (21) give the partial solution $\tau_l^{(p)}$ in terms of elementary functions.

III. SIMPLE REPRESENTATION FOR A_2-COEFFICIENT

In the work [2] (see Appendix C) the coefficient $A_2 \equiv A_2(l)$ included into the physical solution (10), was derived in general but very complicated (integral) form. In particular,

$$ A_2(l) = \frac{1}{\mathcal{P}_2(l)}\left[\frac{(\pi-2)2^{-3(l+2)}}{3\pi(2l-1)(l-2)(l+4)}\mathcal{P}_3(l) - \mathcal{P}_1(l)\right], \quad (29) $$

where

$$ \mathcal{P}_1(l) = \int_0^1 \tau_l^{(p)}(\rho)\frac{\rho^{2l+2}}{(1+\rho^2)^{2l+3}}d\rho, \quad (30) $$

$$\mathcal{P}_2(l) = \int_0^1 v_{4l}(\rho)\frac{\rho^{2l+2}}{(1+\rho^2)^{2l+3}}d\rho = \frac{\sqrt{\pi}\ 2^{-2(l+2)}\Gamma\left(l+3/2\right)}{\Gamma(l/2+3)\Gamma(l/2)}, \tag{31}$$

$$\mathcal{P}_3(l) = -\frac{2^{l+1}}{(l+3)(2l-3)(2l+3)(2l+5)} \times$$
$$\left\{\frac{30}{l+1} - \frac{26}{l+2} + 13 - 4l\left[47 - 2l(2l(l+3)-9)\right] + 2^{l+1}\times\right.$$
$$\left.\left[(l+1)\left[4l(l(4l+3)-17)+45\right]B_{\frac{1}{2}}\left(l+\frac{3}{2},\frac{1}{2}\right) + 8l\left[l(l(4l(l+4)+3)-56)-62\right]B_{\frac{1}{2}}\left(l+\frac{3}{2},\frac{3}{2}\right)\right]\right\}, \tag{32}$$

with the incomplete beta function denoted by $B_z(a,b)$. It is seen that even $\mathcal{P}_3(l)$ can not be called simple. However, the coefficient $\mathcal{P}_1(l)$ can be indeed called very complicated.

The *Mathematica* calculation of the coefficients $A_2(l)$ for any integer $l \geq 3$ shows that

1) for odd values of l the coefficients $A_2(l)$ equal zero;

2) for even values of l the coefficients $A_2(l)$ are reduced to the form

$$A_2(l) = \frac{2-\pi}{\pi^2}\mathcal{A}(l), \tag{33}$$

with $\mathcal{A}(l) = (a_l + \pi b_l)$ where a_l and b_l are rational numbers. Using the effective *Mathematica* code, we have calculated the rational numbers a_l and b_l from $l = 4$ up to $l = 60$ (with step equals 2). Making use of the *Mathematica* operator **FindSequenceFunction** it is possible to find the general simple form of the coefficients a_l and b_l. One should emphasize, that for a given sequence there is a minimal number of terms to enable *Mathematica* to find the formula of the general term. In particular, for the coefficients a_l and b_l these minimal numbers are 10 and 26, corresponding to $l = 4, 6, 8, ...22$ and $l = 4, 6, 8, ...54$, respectively. Thus, application of the *Mathematica* operator **FindSequenceFunction** to the sequences mentioned above yields:

$$\mathcal{A}(l) = \frac{\sqrt{\pi}}{360l(l-2)\Gamma(l+\frac{1}{2})} \times$$
$$\left\{\frac{\left[l(l+1)(688l^4+1376l^3-2480l^2-3168l+465)+450\right]\Gamma\left(\frac{l-1}{2}\right)\Gamma\left(\frac{l+1}{2}\right)}{(2l-3)(2l-1)(2l+1)(2l+3)(2l+5)} - \frac{56}{l-1}\left(\frac{l}{2}!\right)^2\right\}. \tag{34}$$

It is worth noting that inserting Eq.(11) into the rhs of Eq.(10) and using (21), one can represent the physical solution in the form

$$\tau_l(\rho) = \frac{1}{2l+1}u_{4l}(\rho)\mathcal{V}_l(\rho) + \left[B_{l,0} + B_{l,1}(\rho)\right]v_{4l}(\rho), \tag{35}$$

where

$$B_{l,0}=A_2(l)+\frac{8(l-3)!a_{0l}}{15\sqrt{\pi}(2l+1)\Gamma\left(l+\frac{1}{2}\right)}\sum_{m=0}^{l-3}\frac{\Gamma\left(m+\frac{7}{2}\right)\Gamma\left(l-m+\frac{1}{2}\right)(-1)^m}{m!(l-m-3)!}\times \sum_{n=1}^{5}a_{nl}\left(\frac{1}{2m+n}+\frac{1}{2m+n+1}\right), \tag{36}$$

$$B_{l,1}(\rho)=-\frac{8(l-3)!a_{0l}}{15\sqrt{\pi}(2l+1)\Gamma\left(l+\frac{1}{2}\right)}\sum_{m=0}^{l-3}\frac{\Gamma\left(m+\frac{7}{2}\right)\Gamma\left(l-m+\frac{1}{2}\right)(-1)^m}{m!(l-m-3)!}\times \sum_{n=1}^{5}a_{nl}\left(\frac{\rho^{2m+n}}{2m+n}+\frac{\rho^{2m+2n+1}}{2m+n+1}\right). \tag{37}$$

Given that $A_2(l)$ equals zero for odd l, one can derive a simple expression for $B_{l,0}$ with even l. A simple way to solve the problem is to make use of the method, which was applied earlier in order to find the coefficients $\mathcal{A}(l)$. Like the latter coefficients, $B_{l,0}$ have a form $[(2-\pi)/\pi^2](\widetilde{a}_l+\pi\widetilde{b}_l)$, where $\widetilde{a}_l$ and $\widetilde{b}_l$ are rational numbers. Thus, calculating $\widetilde{a}_l$ and $\widetilde{b}_l$ (for even $l\geq 4$) according to Eq.(36), and then using the *Mathematica* operator **FindSequenceFunction**, one finds

$$B_{l,0}=\frac{2-\pi}{360\pi^{3/2}(l-1)}\left[\frac{\sqrt{\pi}\left(40l^4-56l^3+11l^2-53l+30\right)}{2^{l-1}l(l-2)(2l-3)(2l-1)}-\frac{7l\left(\frac{l}{2}-2\right)!\left(\frac{l}{2}-1\right)!}{\Gamma\left(l+\frac{1}{2}\right)}\right]. \tag{38}$$

The minimal length of a sequence enables the Mathematica operator **FindSequenceFunction** to find a simple function that yields the sequences $\widetilde{a}_l$ and $\widetilde{b}_l$ are 10 and 20, corresponding to $l=4,6,8,...22$ and $l=4,6,8,...42$, respectively.

IV. SUBCOMPONENT $\psi_{3,0}^{(2c)}$: SIMPLE REPRESENTATION FOR COEFFICIENT c_l

The last subcomponent we have presented in [2] (see Eq.(98)) was

$$\psi_{3,0}^{(2c)}(\alpha,\theta)=\sum_{l=0}^{\infty}P_l(\cos\theta)(\sin\alpha)^l\phi_l(\rho). \tag{39}$$

It is the physical solution of the individual Fock recurrence relation (see Appendix D [2])

$$\left(\Lambda^2-21\right)\psi_{3,0}^{(2c)}=h_{3,0}^{(2c)} \tag{40}$$

with the rhs of the form

$$h_{3,0}^{(2c)} = \sum_{l=0}^{\infty} P_l(\cos\theta)(\sin\alpha)^l \mathrm{h}_l(\alpha), \tag{41}$$

where

$$\mathrm{h}_l(\alpha) \equiv h_l(\rho) = \frac{2^{1-l}\left(\rho^2+1\right)^{l+\frac{1}{2}}\left[(1-2l)\rho^2+2l+3\right]}{3(2l-1)(2l+3)\rho}. \tag{42}$$

Function $\phi_l(\rho)$ as the solution of the inhomogeneous differential equation

$$\left(1+\rho^2\right)^2\phi_l''(\rho) + 2\rho^{-1}\left[1+\rho^2+l(1-\rho^4)\right]\phi_l'(\rho) + (3-2l)(7+2l)\phi_l(\rho) = -h_l(\rho) \tag{43}$$

was derived in the form (see Eq.(99) [2])

$$\phi_l(\rho) = \phi_l^{(p)}(\rho) + c_l v_{3l}(\rho), \tag{44}$$

where (see Eqs.(100) [2])

$$v_{3l}(\rho) = \left(\rho^2+1\right)^{l-\frac{3}{2}}\left[\frac{(2l-3)(2l-1)}{(2l+3)(2l+5)}\rho^4 + \frac{2(2l-3)}{2l+3}\rho^2 + 1\right]. \tag{45}$$

The particular solution $\phi_l^{(p)}$ of the equation (43) was represented in the form (see Eqs.(101)-(104) [2])

$$\phi_l^{(p)}(\rho) = \frac{2^{-l}\left(\rho^2+1\right)^{l-\frac{3}{2}}}{3(2l-3)(2l-1)(2l+3)(2l+5)}\left[2f_{1l}(\rho) + \frac{2f_{2l}(\rho)+f_{3l}(\rho)}{2l+1}\right], \tag{46}$$

where

$$f_{1l}(\rho) = \left[9-4l(l+2)\right]\rho + \left(13-4l^2\right)\rho^3, \tag{47}$$

$$f_{2l}(\rho) = \left[(2l-3)(2l-1)\rho^4 + 2(2l-3)(2l+5)\rho^2 + (2l+3)(2l+5)\right]\arctan(\rho), \tag{48}$$

$$f_{3l}(\rho) = -\left[(2l+3)(2l+5)\rho^4 + 2(2l-3)(2l+5)\rho^2 + (2l-3)(2l-1)\right] \times \frac{\rho}{l+1}\,{}_2F_1\left(1, l+1; l+2; -\rho^2\right). \tag{49}$$

It was shown (see appendix D [2]) that the coefficient c_l included into solution (44) can be calculated as follows:

$$c_l = \frac{\mathcal{M}_1(l) - \mathcal{M}_2(l)}{\mathcal{M}_3(l)}, \tag{50}$$

where

$$\mathcal{M}_1(l) = \frac{2^{-3l-2}l!\sqrt{\pi}}{3(2l-3)(2l-1)(2l+7)\Gamma(l+3/2)}\,{}_3F_2\left(\frac{2l-1}{4}, \frac{2l+1}{4}, l+1; l+\frac{3}{2}, l+\frac{3}{2}; 1\right), \tag{51}$$

$$\mathcal{M}_3(l) \equiv \int_0^1 v_{3l}(\rho)\frac{\rho^{2l+2}}{\left(\rho^2+1\right)^{2l+3}}d\rho = \frac{2^{-l-\frac{3}{2}}(2l+1)}{(2l+3)(2l+7)}. \tag{52}$$

$$\mathcal{M}_2(l) \equiv \int_0^1 \phi_l^{(p)}(\rho)\frac{\rho^{2l+2}}{\left(\rho^2+1\right)^{2l+3}}d\rho. \tag{53}$$

It is seen that according to Eqs.(50)-(53) the coefficient c_l is represented by very complicated function of l. However, *Mathematica* calculations with any given integer $l \geq 0$ show that the parameter c_l has a form $c_{0,l} + c_{1,l}\pi + c_{2,l}\ln 2$ where $c_{i,l}$ $(i = 0, 1, 2)$ are rational numbers. Using the Mathematica operator **FindSequenceFunction**, one obtains the following simple result

$$c_l = \frac{2l+1-(\pi/2)-\Phi(-1,1,l+1)}{3(2l-3)(2l-1)(2l+1)2^l}, \tag{54}$$

where $\Phi(z, s, a)$ is the Lerch transcendent. Note that for the case under consideration we have

$$\Phi(-1,1,l+1) = \frac{1}{2}\left[\psi\left(\frac{l}{2}+1\right) - \psi\left(\frac{l}{2}+\frac{1}{2}\right)\right] = \frac{1}{2}\left(H_{\frac{l}{2}} - H_{\frac{l-1}{2}}\right), \tag{55}$$

where $\psi(z)$ and H_z are the digamma function and harmonic number, respectively.

The minimal length of a sequence enables the *Mathematica* to find a simple function that yields the sequence $c_{0,l}$ is 22, whereas for $c_{1,l}$ and $c_{2,l}$ it equals 6.

Warning: There is the following (software) **bug** in the *Mathematica* version 10.3.

The action of the *Mathematica* operators **FullSimplify** or **FunctionExpand** on the generalized hypergeometric function

$${}_3F_2\left(\frac{2l-1}{4}, \frac{2l+1}{4}, l+1; l+\frac{3}{2}, l+\frac{3}{2}; 1\right)$$

with any given integer l produces a meaningless expression. One should emphasize that the previous *Mathematica* versions, e.g., 10.0 or 9.0 give the correct results. For example, for $l = 2$, one obtains:

$l := 2;$

$$\textbf{FunctionExpand}\left[HypergeometricPFQ\left[\left\{\frac{2l-1}{4}, \frac{2l+1}{4}, l+1\right\}, \left\{l+\frac{3}{2}, l+\frac{3}{2}\right\}, 1\right]\right] = \frac{2}{49}(-352+254\sqrt{2}).$$

Notice that the considered generalized hypergeometric function is included into Eq.(51).

---

[1] V. A. Fock, "On the Schrödinger Equation of the Helium Atom", Izv. Akad. Nauk SSSR, Ser. Fiz. **18**, 161-174 (1954).

[2] E. Z. Liverts and N. Barnea , "Angular Fock coefficients: Refinement and further development", Phys. Rev. A **92**, 042512 (2015).

[3] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, "Integrals and Series. Vol 3. More Special Functions", Gordon and Breach S. P., New York, 1986.

[4] http://functions.wolfram.com/HypergeometricFunctions/Hypergeometric2F1/17/01/03/0005/

[5] http://functions.wolfram.com/HypergeometricFunctions/Hypergeometric2F1/16/01/01/0002/

[6] http://functions.wolfram.com/HypergeometricFunctions/Hypergeometric2F1/03/06/07/07/0001/

[7] http://functions.wolfram.com/HypergeometricFunctions/Hypergeometric2F1/21/01/02/01/0003/

[8] http://functions.wolfram.com/HypergeometricFunctions/Hypergeometric3F2/03/07/02/0001/

[9] http://functions.wolfram.com/HypergeometricFunctions/Hypergeometric2F1/17/01/03/0006/